

# General Agriculture Application Information

For

**GRASS (PASTURES, HAY and LAWNS)**  
Soil and Foliar Applications

## Products:

**The Bio-N-Liven Answer®**

**The Carbon Answer®**

**The Mineral Electrolyte Answer®**

**Note:** Application rates are “general.”  
They have been compiled over the previous 15 years  
from field tests and long-term usage by growers.

Such tests are ongoing and this table will  
be updated as more information  
is made available.



Revised January 2005  
[www.ECandS.net](http://www.ECandS.net)



# THE ANSWER® 2005 PRODUCT APPLICATION RATES

## GRASS (PASTURES, HAY and LAWNS)

Application Area	The Answer® Products	Application Rate	Application Time	Dilution Rate	Method of Application	Field Observations
Soil	Bio-N-Liven Answer®	Fertile Soil: (24 fl. Oz./acre) Moderate Soil: (32 fl. Oz./acre) Problem Soil: (40 fl. Oz./acre)	Established pastures and lawns: Early Spring New lawns: Before planting	Aerial - 1:20, 1:50 Ground - at least 1:100 Dilution rate may vary depending on the spray equipment used.	Broadcast: Aerial or ground. Spray diluted product evenly on soil using any conventional spray equipment. New crops: Incorporate product into soil. Established crops: Product must be applied when soil is moist and rain is expected in 3-5 days.	INCREASED: Soil fertility, water holding capacity, drainage, organic matter content, neutralization of soil pH
Soil	Mineral Electrolyte Answer®	Fertile Soil: (6 fl. Oz./acre) Moderate Soil: (8 fl.oz./acre) Problem Soil: (10 fl. Oz./acre)	Established pastures and lawns: Early Spring New lawns: Before planting	Aerial - 1:20, 1:50 Ground - at least 1:100 Dilution rate may vary depending on the spray equipment used.	Broadcast: Aerial or ground. Spray diluted product evenly on soil using any conventional spray equipment. New crops: Incorporate product into soil. Established crops: Product must be applied when soil is moist and rain is expected in 3-5 days.	INCREASED: Soil fertility, water holding capacity, drainage, organic matter content, neutralization of soil pH
Soil	Carbon Answer®	Fertile Soil: (2 fl. Oz./acre) Moderate Soil: (2 fl. Oz./acre) Problem Soil: (2 fl. Oz./acre)	Established pastures and lawns: Early Spring New lawns: Before planting	Aerial - 1:20, 1:50 Ground - at least 1:100 Dilution rate may vary depending on the spray equipment used.	Broadcast: Aerial or ground. Spray diluted product evenly on soil using any conventional spray equipment. New crops: Incorporate product into soil. Established crops: Product must be applied when soil is moist and rain is expected in 3-5 days.	INCREASED: Soil fertility, water holding capacity, drainage, organic matter content, neutralization of soil pH

**APPLICATION COLOR GUIDE**





# THE ANSWER® 2005 PRODUCT APPLICATION RATES

## GRASS (PASTURES, HAY and LAWNS)

Application Area	The Answer® Products	Application Rate	Application Time	Dilution Rate	Method of Application	Field Observations
Foliar	Bio-N-Liven Answer®	(12 fl. Oz./acre)	Twice: First growth in Spring and in 4-6 weeks	Aerial - 1:20, 1:50 Ground - at least 1:100 Dilution rate may vary depending on the spray equipment used.	Broadcast: Aerial or ground. Spray evenly on soil using any conventional spray equipment. For best uptake, apply in evening or early morning. Heavy rain or aerial irrigation within 24 hours of application will diminish product effectiveness.	IMPROVED: Vigor of plant, regrowth, greener and thicker leaves, yield, protein content
Foliar	Mineral Electrolyte Answer®	(3 fl. Oz./acre)	Twice: First growth in Spring and in 4-6 weeks	Aerial - 1:20, 1:50 Ground - at least 1:100 Dilution rate may vary depending on the spray equipment used.	Broadcast: Aerial or ground. Spray evenly on soil using any conventional spray equipment. For best uptake, apply in evening or early morning. Heavy rain or aerial irrigation within 24 hours of application will diminish product effectiveness.	IMPROVED: Vigor of plant, regrowth, greener and thicker leaves, yield, protein content
Foliar	Carbon Answer®	(2 fl. Oz./acre)	Twice: First growth in Spring and in 4-6 weeks	Aerial - 1:20, 1:50 Ground - at least 1:100 Dilution rate may vary depending on the spray equipment used.	Broadcast: Aerial or ground. Spray evenly on soil using any conventional spray equipment. For best uptake, apply in evening or early morning. Heavy rain or aerial irrigation within 24 hours of application will diminish product effectiveness.	IMPROVED: Vigor of plant, regrowth, greener and thicker leaves, yield, protein content

**APPLICATION COLOR GUIDE**



SOIL



SEED



FOLIAR